


**Instructions for Use**  
**ENSTILAR<sup>®</sup>**  
**(calcipotriene and betamethasone dipropionate)**  
**Foam, 0.005%/0.064%**

Read the Patient Information and Instructions for Use before you use Enstilar<sup>®</sup> Foam.

**Important information:** For skin use only (topical use). Do not get Enstilar<sup>®</sup> Foam in your mouth, eyes or vagina. If you accidentally get Enstilar<sup>®</sup> Foam on the face, in the mouth or in the eyes, wash the area with water right away. Do not swallow Enstilar<sup>®</sup> Foam.


**How to apply ENSTILAR<sup>®</sup> Foam:**

Follow your doctor's instructions on how much Enstilar<sup>®</sup> Foam to use and where to use it.



**Wash your hands before applying Enstilar<sup>®</sup> Foam.**

**Step 1:** Remove the cap from the can. Shake the can before use.



**Step 2:** Hold the can at least 1.5 inches from the affected area.

	<p><b>Step 3:</b> The foam can be sprayed holding the can in any position except horizontally.</p>
	<p><b>Step 4:</b> Gently rub in Enstilar<sup>®</sup> Foam into your affected skin areas.</p>
<p><b>Step 5:</b> After applying Enstilar<sup>®</sup> Foam, put the cap back on the can.</p> <p><b>Step 6:</b> Wash your hands after using Enstilar<sup>®</sup> Foam unless you are using the medicine to treat your hands.</p>	

This Instructions for Use has been approved by the U.S. Food and Drug Administration.

Manufactured by:  
LEO Laboratories Ltd.  
285 Cashel Road  
Dublin 12, Ireland

or

Colep Laupheim GmbH & Co. KG  
Fockestraße 12  
88471 Laupheim  
Germany (DE)

Distributed by:  
LEO Pharma Inc.  
Seven Giralda Farms  
Madison, NJ 07940, USA

Issued: 06/2017